

Decimal Fraction

1. It is a fraction in which denominators are powers of 10.

$$1 / 10 = 1 \text{ tenth} = .1$$

$$1/100 = 1 \text{ hundredth} = .01$$

$$99/100 = 99 \text{ hundredths} = .99$$

$$7 / 1000 = 7 \text{ thousandths} = .007, \text{ etc}$$

2. Conversion of a Decimal into Vulgar Fraction

Put 1 in the denominator under the decimal point and annex with it as many zeros as is the number of digits after the decimal point. Now, remove the decimal point and reduce the fraction to its lowest terms.

$$\text{Thus, } 0.25 = 25/100=1/4$$

$$2.008= 2008/1000 =251/125$$

3. Annexing Zeros and Removing Decimal Signs

Annexing zeros to the extreme right of a decimal fraction does not change its value. Thus, $0.8 = 0.80 = 0.800$, etc.

If numerator and denominator of a fraction contain the same number of decimal places, then we remove the decimal sign. Thus, $1.84/2.99=184/299=8/13$

4. Operations on Decimal Fractions

i. Addition and Subtraction of Decimal Fractions: The given numbers are so placed under each other that the decimal points lie in one column. The numbers so arranged can now be added or subtracted in the usual way.

ii. Multiplication of a Decimal Fraction By a Power of 10: Shift the decimal point to the right by as many places as is the power of 10.

$$\text{Thus, } 5.9632 \times 100 = 596.32; 0.073 \times 10000 = 730.$$

iii. Multiplication of Decimal Fractions: Multiply the given numbers

considering them without decimal point. Now, in the product, the decimal point is marked off to obtain as many places of decimal as is the sum of the number of decimal places in the given numbers. Suppose we have to find the product ($.2 \times 0.02 \times .002$).

$$\text{Now, } 2 \times 2 \times 2 = 8. \text{ Sum of decimal places} = (1 + 2 + 3) = 6.$$

$$.2 \times .02 \times .002 = .000008$$

iv. Dividing a Decimal Fraction By a Counting Number

Divide the given number without considering the decimal point, by the given counting number. Now, in the quotient, put the decimal point to give as many places of decimal as there are in the dividend.

$$\text{Suppose we have to find the quotient } (0.0204 \div 17). \text{ Now, } 204 \div 17 = 12.$$

$$\text{Dividend contains 4 places of decimal. So, } 0.0204 \div 17 = 0.0012$$

v. Dividing a Decimal Fraction By a Decimal Fraction

Multiply both the dividend and the divisor by a suitable power of 10 to make divisor a whole number. Now, proceed as above.

$$0.0066/0.11= (0.00066*100) / 0.11 * 100= 0.066/11=0.06$$

5. Comparison of Fractions

Suppose some fractions are to be arranged in ascending or descending order of magnitude, then convert each one of the given fractions in the decimal form, and arrange them accordingly.

Let us to arrange the fractions $\frac{3}{5}$, $\frac{6}{7}$ and $\frac{7}{9}$ in descending order.

Now,

$$\frac{3}{5} = 0.6, \frac{6}{7} = 0.857, \frac{7}{9} = 0.777\dots$$

Since, $0.857 > 0.777\dots > 0.6$. $\frac{6}{7} > \frac{7}{9} > \frac{3}{5}$

6. Recurring Decimal

If in a decimal fraction, a figure or a set of figures is repeated continuously, then such a number is called a recurring decimal. In a recurring decimal, if a single figure is repeated, then it is expressed by putting a dot on it. If a set of figures is repeated, it is expressed by putting a bar on the set. Thus, $\frac{1}{3} = 0.333\dots = 0.\dot{3}$; $\frac{22}{7} = 3.142857142857\dots = 3.14285\bar{7}$.

Pure Recurring Decimal

A decimal fraction, in which all the figures after the decimal point are repeated, is called a pure recurring decimal. Converting a Pure Recurring Decimal into Vulgar Fraction: Write the repeated figures only once in the numerator and take as many nines in the denominator as is the number of repeating figures.

Thus, $0.5 = \frac{5}{9}$, $0.53 = \frac{53}{99}$; $0.067 = \frac{67}{999}$, etc.

Mixed Recurring Decimal

A decimal fraction in which some figures do not repeat and some of them are repeated, is called a mixed recurring decimal.

Eg. $0.1733333\dots = 0.17\dot{3}$.

Converting a Mixed Recurring Decimal Into Vulgar Fraction

In the numerator, take the difference between the number formed by all the digits after decimal point (taking repeated digits only once) and that formed by the digits which are not repeated. In the denominator, take the number formed by as many nines as there are repeating digits followed by as many zeros as is the number of non-repeating digits.

Thus, $0.16 = \frac{16 - 1}{90} = \frac{15}{90} = \frac{1}{6}$, $0.227\dot{3} = \frac{2273 - 22}{9900} = \frac{2251}{9900}$

Problems with solutions

1. What decimal of an hour is a second ?

Solution

$$\text{Required decimal} = \frac{1}{60 \times 60} = \frac{1}{3600} = .00027$$

2. If $2994 \div 14.5 = 172$, then $29.94 \div 1.45 = ?$

Solution

$$\begin{aligned} \frac{29.94}{1.45} &= \frac{299.4}{14.5} \\ &= \left(\frac{2994}{14.5} \times \frac{1}{10} \right) \text{ [Here, Substitute 172 in the place of } 2994/14.5 \text{]} \\ &= \frac{172}{10} \\ &= 17.2 \end{aligned}$$

3. When 0.232323..... is converted into a fraction, then the result is:

Solution

$$0.232323... = \frac{23}{99}$$

4. The expression $(11.98 \times 11.98 + 11.98 \times x + 0.02 \times 0.02)$ will be a perfect square for x equal to:

Solution

$$\text{Given expression} = (11.98)^2 + (0.02)^2 + 11.98 \times x.$$

For the given expression to be a perfect square, we must have

$$11.98 \times x = 2 \times 11.98 \times 0.02 \text{ or } x = 0.04$$

5. If $\frac{144}{0.144} = \frac{14.4}{x}$, then the value of x is:

Solution

$$\begin{aligned} \frac{144}{0.144} &= \frac{14.4}{x} \\ \frac{144 \times 1000}{144} &= \frac{14.4}{x} \\ x &= \frac{14.4}{1000} = 0.0144 \end{aligned}$$